

HULTMAN ELEMENTS FOR THE HYPEROCTAHEDRAL GROUPS

ALEXANDER WOO

ABSTRACT. Hultman, Linusson, Shareshian, and Sjöstrand gave a pattern avoidance characterization of the permutations for which the number of chambers of its associated inversion arrangement is the same as the size of its lower interval in Bruhat order. Hultman later gave a characterization, valid for an arbitrary finite reflection group, in terms of distances in the Bruhat graph. On the other hand, the pattern avoidance criterion for permutations had earlier appeared in independent work of Sjöstrand and Gasharov and Reiner. We give characterizations of the elements of the hyperoctahedral groups satisfying Hultman's criterion that is in the spirit of those of Sjöstrand and of Gasharov and Reiner. We also give a pattern avoidance criterion using the notion of pattern avoidance defined by Billey and Postnikov.

1. INTRODUCTION

Let $w \in S_n$ be a permutation, and let $H_{i,j}$ denote the hyperplane in \mathbb{R}^n defined by $x_i = x_j$. The **inversion arrangement** of w is the hyperplane arrangement in \mathbb{R}^n given by

$$\mathcal{A}_w := \{H_{i,j} \mid i < j, w(i) > w(j)\}.$$

Any hyperplane arrangement \mathcal{A} cuts \mathbb{R}^n into a number of **chambers**, which are defined as the connected components of $\mathbb{R}^n \setminus \mathcal{A}$. Hence we can associate an invariant $c(w)$, the number of chambers of \mathcal{A}_w , to any permutation w .

On the other hand, if we let $[\text{id}, w]$ denote the interval from the identity to w in Bruhat order, we can also associate to w the invariant $s(w) := \#[\text{id}, w]$. Following a conjecture of Postnikov [21, Remark 24.2], Hultman, Linusson, Shareshian, and Sjöstrand [15] proved that $c(w) \leq s(w)$ for any permutation w , and, furthermore, that $c(w) = s(w)$ if and only if w avoids the permutations 4231, 35142, 42513, and 351624.

The concepts defined above make sense for an arbitrary finite reflection group W . Given any element $w \in W$, one can define an inversion arrangement \mathcal{A}_w and an invariant $c(w)$ counting its chambers. There is also a notion of Bruhat order, so one can associate to w an invariant $s(w)$. Indeed, Hultman, Linusson, Shareshian, and Sjöstrand [15] actually showed that $c(w) \leq s(w)$ for elements w of any finite reflection group W , and, in slightly later work, Hultman [16] gave a characterization of the elements w for which $c(w) = s(w)$ in terms of certain conditions on distances in the Bruhat graph of W . We will call the elements $w \in W$ for which $c(w) = s(w)$ **Hultman elements**.

The set of permutations avoiding 4231, 35142, 42513, and 351624 had previously appeared in two independent but related places in the literature. A permutation is said

Date: May 22, 2018.

to be **defined by inclusions** if the interval $[\text{id}, w]$ is defined by inclusion conditions. To be more precise, this means there are (possibly empty) sets $B(w)$ and $T(w)$ such that $u \in [\text{id}, w]$ if and only if $\{u(1), \dots, u(q)\} \subseteq \{1, \dots, p\}$ for all $(p, q) \in B(w)$ and $\{1, \dots, p\} \subseteq \{u(1), \dots, u(q)\}$ for all $(p, q) \in T(w)$. Gasharov and Reiner [12] showed that a permutation is defined by inclusions if and only if it avoids 4231, 35142, 42513, and 351624. The **right hull** of a permutation w is the rectilinear convex hull of its graph and the points $(1, 1)$ and (n, n) . A permutation w satisfies the **right hull condition** if $u \leq w$ for every permutation u whose graph fits in the right hull of w . Sjöstrand [25] showed that the same set of permutations are the ones satisfying the right hull condition.

Our goal in this paper is to give a characterization of Hultman elements for the hyperoctahedral groups B_n that is in the spirit of the Gasharov–Reiner and Sjöstrand characterizations for S_n , considering B_n as a subgroup of S_{2n} . We say that an element $w \in B_n \subseteq S_{2n}$ is **defined by pseudo-inclusions** if it is defined by inclusions (as an element of S_{2n}), possibly with the additional condition $\#(\{u(1), \dots, u(n)\} \cap \{1, \dots, n\}) = n - 1$. One can similarly define a relaxation of the right hull condition. Furthermore, there is a generalization of pattern avoidance to arbitrary Coxeter groups due to Billey and Postnikov [5], which we call **BP avoidance**. Our main theorem is as follows. (Precise definitions are given in Section 2.)

Main Theorem. *Let $w \in B_n$. Then the following are equivalent.*

- (1) *The number of chambers of the inversion arrangement \mathcal{A}_w is equal to the number of elements in $[\text{id}, w]$.*
- (2) *For any $u \leq w$, the directed distance from u to w in the Bruhat graph is the same as the undirected distance from u to w .*
- (3) *The element w is defined by pseudo-inclusions.*
- (4) *The element w satisfies the relaxed right hull condition.*
- (5) *The element w BP avoids $4231 \in S_4$, $35142, 42513 \in S_5$, $351624 \in S_6$, $563412, 653421, 645231, 635241, 624351, 642531, 536142, 426153, 462513, 623451 \in B_3$, $47618325, 46718235, 57163824, 37581426, 47163825, 46172835, 37518426, 35718246, 37145826, 37154826, 52618374, 42681375, 42618375, 35172846 \in B_4$, and $3517294a68, 3517924a68, 3617294a58 \in B_5$.*

The equivalence of the first 2 statements is due to Hultman. Our proof showing the equivalence of the second condition with the remainder largely follows the proof of Hultman [16] recovering the pattern avoidance condition due to Hultman, Linusson, Shareshian and Sjöstrand [15] from the distance condition, though a number of details are significantly more complicated.

First we give a proof of the equivalence of the third and fourth conditions; this direct proof of equivalence is new even for S_n . Afterwards, given an element $w \in B_n$ satisfying the relaxed right hull condition, we give a more complicated variant of Hultman’s proof showing that the distance condition is satisfied in the Bruhat graph.

Second, given an element $w \in B_n$ not defined by pseudo-inclusions, we show that w must BP contain an element of S_m or B_m not defined by pseudo-inclusions for some $m \leq 5$. A computer calculation then shows that the above list is the minimal possible. Another

calculation shows that none of the elements on the list satisfy the distance condition. We finally prove that, if w BP contains u and u fails to satisfy the distance condition, then w must also fail to satisfy the distance condition. It is possible to avoid the first calculation by a more intricate but quite tedious version of the argument of Gasharov and Reiner.

One can replace BP avoidance with avoidance of signed permutations at the cost of significantly lengthening the list of patterns to be avoided. We will see, however, that BP avoidance is the natural notion to use in this context.

This work has several natural possible extensions. First, it would be interesting to extend this characterization to all finite reflection groups. An extension to type D using the methods of this paper is likely possible and may be the subject of a future paper. An underlying principle guides the expected extension. Fulton [11] defined the notion of the **essential set** $E(w)$ of a permutation w , which is a set of conditions that characterizes when $u \in [\text{id}, w]$. Given $w \in S_n$, one can associate to each element of $(p, q) \in E(w)$ a permutation $v(p, q, r_w(p, q))$ such that u fails the condition specified by (p, q, r) if and only if $v \geq u(p, q, r)$. Given an element $w \in B_n \subseteq S_{2n}$, we can similarly associate to each pair

$$\{(p, q, r), (2n + 2 - p, 2n - q, p - q - 1 + r)\} \subseteq E(w)$$

an element

$$v(p, q, r) = v(2n + 2 - p, 2n - q, p - q - 1 + r) \in B_n$$

such that $u \in B_n$ fails both conditions if and only if $u \geq v(p, q, r)$. It turns out that, for both S_n and B_n , an element w is Hultman if and only if $v(p, q, r)$ has only one reduced expression for all $(p, q, r) \in E(w)$. Unfortunately, this condition cannot be stated in terms of the Coxeter-theoretic definition of coessential set given by Reiner, Yong, and the author in [23] because some conditions in $E(w)$ can be implied by other conditions in $E(w)$ when only considering elements of B_n . We discuss this in Section 4 using recent work of Anderson [2]. Given the nature of Hultman's proof and ours, we expect the recent work of Gobet [14] on analogues of cycle decompositions for finite reflection groups to be useful in any general proof of a general statement for all finite reflection groups.

Furthermore, several additional results concerning the permutations defined by inclusions have potential analogues in B_n . Lewis and Morales [20] showed that, if w is a permutation defined by inclusions, then the number of invertible matrices over \mathbb{F}_q supported on the complement of the diagram of w is related to the rank generating function of the interval $[\text{id}, w]$ and hence is a q -analogue of $s(w)$, but no analogous result for B_n has been found. Gasharov and Reiner [12] gave a presentation of the cohomology ring $H^*(X_w)$ for a Schubert variety corresponding to a permutation $w \in S_n$ defined by inclusions; indeed they were originally interested only in the case where X_w is smooth and defined the notion of a permutation defined by inclusions because their results naturally generalized to this case. It would be interesting to extend their results to Schubert varieties associated to Hultman elements of B_n . It is likely that any presentation of the cohomology ring would involve the theta polynomials of Buch, Kresch, and Tamvakis [7]. Also, Ulfarsson and the author showed that the Kostant–Kumar polynomials for permutations defined by inclusions are products of not necessarily distinct roots [26, Cor. 6.6]. It would be interesting to prove the converse as well as extend this result to B_n . Finally, Albert and Brignall [1] computed the generating function counting permutations defined

by inclusions; one possible path to an analogous result for B_n would be to understand the proof of Albert and Brignall in the context of the staircase diagrams of Richmond and Slofstra [24].

We organize the paper as follows. Section 2 gives various preliminaries, including some details of how BP pattern avoidance and cycle decomposition work for B_n . The details on BP avoidance for B_n do not seem to have previously appeared in print. Section 3 contains the proof of our main theorem. Section 4 remarks on the relationship between our main theorem and the Coxeter-theoretic coessential set.

2. PRELIMINARIES

2.1. Bruhat order and inversion arrangements for permutations. A **transposition** is a permutation of the form $t_{i,j} := (i\ j)$, swapping two elements and fixing all the others. A **simple transposition** is a transposition $s_i := t_{i,i+1}$. Let $w \in S_n$ be a permutation. The **length** of w , denoted $\ell(w)$, is the minimum number such that w is a product of $\ell(w)$ simple transpositions. A pair (i, j) with $1 \leq i < j \leq n$ is an **inversion** of w if $w(i) > w(j)$. It turns out that $\ell(w)$ is the number of inversions of w . The **absolute length** of w , denoted $\ell_T(w)$, is the minimum number such that w is a product of $\ell_T(w)$ (not necessarily simple) transpositions. Let $\text{cyc}(w)$ be the number of cycles in the cycle decomposition of w . Then $\ell_T(w) = n - \text{cyc}(w)$ for $w \in S_n$.

The symmetric group S_n has a partial order known as **Bruhat order**. It can be defined as the transitive closure of the covering relation where $u \prec w$ if $w = ut_{i,j}$ for some transposition $t_{i,j}$ and $\ell(w) = \ell(u) + 1$. Alternatively, Bruhat order can also be defined by the **tableau criterion**. Given p, q with $1 \leq p, q \leq n$, define

$$r_w(p, q) := \#\{k \mid 1 \leq k \leq q, p \leq w(k) \leq n\}.$$

The function r_w is called the **SW rank function** of w . Then $u \leq w$ if and only if $r_u(p, q) \leq r_w(p, q)$ for all p, q . We let $s(w)$ be the number of elements in the interval $[\text{id}, w]$.

Example 2.1. Consider $w = 35142$ and $u = 13254$. (All permutations in this paper are written in one line notation, except that $(i\ j)$ denotes the transposition switching i and j .) We have $u < w$ since

$$u = 13254 \prec 31254 \prec 32154 \prec 35124 \prec 35142 = w.$$

On the other hand, the rank functions r_u and r_w , displayed with values in a matrix, are

$$r_u = \begin{array}{ccccc} 1 & 2 & 3 & 4 & 5 \\ 0 & 1 & 2 & 3 & 4 \\ 0 & 1 & 1 & 2 & 3 \\ 0 & 0 & 0 & 1 & 2 \\ 0 & 0 & 0 & 1 & 1 \end{array} \quad \text{and} \quad r_w = \begin{array}{ccccc} 1 & 2 & 3 & 4 & 5 \\ 1 & 2 & 2 & 3 & 4 \\ 1 & 2 & 2 & 3 & 3 \\ 0 & 1 & 1 & 2 & 2 \\ 0 & 1 & 1 & 1 & 1 \end{array}.$$

We see that every entry in r_u is smaller than the corresponding entry of r_w .

The **inversion arrangement** of w is the hyperplane arrangement \mathcal{A}_w in \mathbb{R}^n consisting of the hyperplanes defined by $x_i - x_j = 0$ for all inversions (i, j) of w . The **chambers**

of a hyperplane arrangement \mathcal{A} are the connected components of $\mathbb{R}^n \setminus (\bigcup_{H \in \mathcal{A}} H)$. We let $c(w)$ denote the number of chambers of \mathcal{A}_w .

Example 2.2. For $w = w_0$, where w_0 is the permutation with $w_0(i) := n + 1 - i$ for all i , the arrangement \mathcal{A}_{w_0} is the full braid arrangement for S_n , and $c(w) = n!$. For $w = \text{id}$, \mathcal{A}_{id} is the empty arrangement. If $w = 3412 \in S_4$, then \mathcal{A}_w consists of the hyperplanes defined by $x_4 - x_1$, $x_3 - x_2$, $x_3 - x_1$, and $x_4 - x_2$, and $c(3412) = 14$. If $w = 4231 \in S_4$, then \mathcal{A}_w consists of the hyperplanes defined by $x_4 - x_1$, $x_4 - x_2$, $x_4 - x_3$, $x_3 - x_1$, and $x_2 - x_1$, and $c(4231) = 18$.

Given permutations $v \in S_m$ and $w \in S_n$ with $m \leq n$, we say that w (**pattern**) **contains** v if there exist indices $1 \leq i_1 < \dots < i_m \leq n$ such that, for all j, k with $1 \leq j < k \leq m$, $v(j) > v(k)$ if and only if $w(i_j) > w(i_k)$. For example, $w = 48631725$ contains $v = 35142$ using indices $i_1 = 1, i_2 = 2, i_3 = 5, i_4 = 6$, and $i_5 = 7$. We say w (**pattern**) **avoids** v if w does not contain v . For example, $w = 68435271$ avoids $v = 35142$.

Hultman, Linusson, Shareshian, and Sjöstrand [15] proved the following.

Theorem 2.1. *Let $w \in S_n$ be a permutation. Then*

- (1) $c(w) \leq s(w)$.
- (2) $c(w) = s(w)$ if and only if w avoids $4231, 35142, 42513$, and 351624 .

Example 2.3. For $w = w_0$, the interval $[\text{id}, w_0]$ is all of S_n , so $c(w) = s(w) = n!$. For $w = 3412$, we have $c(w) = s(w) = 14$. On the other hand, for $w = 4231$, we have $c(w) = 18$, but $s(w) = 20$.

2.2. Permutations defined by inclusions and the right hull property. Given a permutation w , the **diagram** for w , denoted $D(w)$, is defined as follows. Draw a grid of $n \times n$ boxes, mark the entries of the permutation w as in its permutation matrix, and cross out the boxes directly above and directly to the right of a permutation entry. The remaining boxes are the diagram. This means

$$D(w) = \{(p, q) \mid p > w(q), w^{-1}(p) > q\}.$$

Note that $(p, q) \in D(w)$ if and only if $(q, w^{-1}(p))$ is not an inversion.

The **coessential set** $E(w)$ is the set of northeast-most boxes in a connected component of $D(w)$. More precisely, $(p, q) \in E(w)$ if $(p, q) \in D(w)$, $(p-1, q) \notin D(w)$, and $(p, q+1) \notin D(w)$. Alternatively,

$$E(w) = \{(p, q) \mid w^{-1}(p-1) \leq q < w^{-1}(p), w(q) < p \leq w(q+1)\}.$$

It is a lemma of Fulton [11] that $u \leq w$ if $r_u(p, q) \leq r_w(p, q)$ for all $(p, q) \in E(w)$, and, furthermore, $E(w)$ is the unique minimal set that determines the Bruhat interval $[\text{id}, w]$. (To be precise, Fulton defined an **essential set** using a NW rank function and proved a lemma about the essential set that is equivalent to the one stated here.)

Example 2.4. (This is copied from [26, Example 2.2].) Let $w = 819372564$. Then the diagram and coessential set of w are as in Figure 1. In particular,

$$E(w) = \{(2, 2), (4, 4), (4, 6), (6, 7), (9, 2)\},$$

with $r_w(2, 2) = 1, r_w(4, 4) = 2, r_w(4, 6) = 3, r_w(6, 7) = 3$, and $r_w(9, 2) = 0$.




FIGURE 1. Diagram and essential set for $w = 819372564$.

A permutation w is **defined by inclusions** if, for all $(p, q) \in E(w)$, we have $r_w(p, q) = \max(0, q - p + 1) = r_{\text{id}}(p, q)$. Note that $r_w(p, q) \geq \max(0, q - p + 1)$ for all p, q for any permutation w , so being defined by inclusions means that the rank numbers are as small as possible for elements of the coessential set. We use this terminology because $r_u(p, q) \leq 0$ if and only if $\{u(1), \dots, u(q)\} \subseteq \{1, \dots, p - 1\}$ and $r_u(p, q) \leq q - p + 1$ if and only if $\{1, \dots, p - 1\} \subseteq \{u(1), \dots, u(q)\}$. (The first statement is obvious from the definitions. For the second, see Lemma 3.1.) Alternatively, w is defined by inclusions if the Schubert variety X_w is defined by conditions of the form $F_q \subseteq E_{p-1}$ (when $r_w(p, q) = 0$) and the form $E_{p-1} \subseteq F_q$ (when $r_w(p, q) = q - p + 1$).

Gasharov and Reiner [12] showed that a permutation w is defined by inclusions if and only if w avoids 4231, 35142, 42513, and 351624.

Example 2.5. The permutation $w = 819372564$ in Example 2.4 is not defined by inclusions since $(4, 4) \in E(w)$ and $r_w(4, 4) = 2 \neq 4 - 4 + 1$. We see that w contains 4231 (in several ways).

Given a permutation w , define the **right hull** of w to be the set $H(w)$ of points (i, j) satisfying *both* of the following:

- $1 \leq i \leq w(k)$ for some $k \leq j$
- $w(k) \leq i \leq n$ for some $k \geq j$.

We say that a permutation v is in $H(w)$ and write $v \subseteq H(w)$ if $(v(j), j) \in H(w)$ for all j , $1 \leq j \leq n$. In other words, for this purpose, we think of a permutation v as the set of points $\{(v(j), j) \mid 1 \leq j \leq n\}$.

Given any set $S \subseteq \mathbb{R}^2$ (which we index with matrix coordinates for the purposes of this paper), let the **rectilinear SW-NE hull** of S be the set of points

$$H(S) := \{(p, q) \mid p_1 \geq p \geq p_2, q_1 \leq q \leq q_2 \text{ for some } (p_1, q_1) \text{ and } (p_2, q_2) \text{ in } S\}.$$

The right hull of w can be thought of as the rectilinear SW-NE hull of w .

Note that, if $u \leq w$, then $u \subseteq H(w)$. A permutation w satisfies the **right hull condition** if the converse is true, meaning that, for all $u \subseteq H(w)$, we have $u \leq w$.

Sjöstrand [25] showed that a permutation w satisfies the right hull condition if and only if w avoids 4231, 35142, 42513, and 351624. He asked for an explanation of the connection between this result and the result of Gasharov and Reiner, which we will provide in Section 3.1.

Example 2.6. Let $w = 819372564$ as in Example 2.4. The right hull $H(w)$ is the unshaded region in Figure 2. If $u = 168523479$, then $u \in H(w)$, but $u \not\leq w$ since $r_u(4,4) = 3$ but $r_w(4,4) = 2$. Hence w does not satisfy the right hull condition.




FIGURE 2. Right hull for $w = 819372564$.

2.3. Inversion arrangements for finite Coxeter groups. Let (W, S) be a **Coxeter group**. This is a group along with a distinguished set of generators S such that the defining relations are $s^2 = \text{id}$ for all $s \in S$ and $(st)^{m(s,t)} = \text{id}$ for all pairs $s, t \in S$, for some $m(s, t) \in \{2, 3, \dots\} \cup \{\infty\}$. The symmetric group S_n is a Coxeter group with generators $s_i = (i \ i+1)$. In this case, we have $m(s_i, s_j) = 2$ if $|j - i| \geq 2$ and $m(s_i, s_{i+1}) = 3$. A **Coxeter group isomorphism** $\phi : (W', S') \rightarrow (W, S)$ is a group isomorphism $\phi : W' \rightarrow W$ that induces a bijection between S' and S . For a general introduction to the combinatorics of Coxeter groups, see [6].

Given a Coxeter group (W, S) , a **reflection** is a conjugate of an element of S ; we denote the set of all reflections by T . Given $w \in W$, the **length** of w , denoted $\ell(w)$, is the minimum number of elements of S whose product is w , and the **absolute length** of w , denoted $\ell_T(w)$, is the minimum number of elements of T whose product is w . **Bruhat order** is the transitive closure of the covering relation defined by $v \prec w$ if $\ell(w) = \ell(v) + 1$ and $w = vt$ for some $t \in T$. As before, we let $s(w)$ denote the number of elements in the interval $[\text{id}, w]$. The **inversion set** of w , denoted $\text{Inv}(w)$, is the set

$$\text{Inv}(w) := \{t \in T \mid wt < w\},$$

where $<$ denotes Bruhat order. (Hultman [16] gives a definition of inversion equivalent to the condition that $tw < w$. This forces him to multiply permutations backwards, which

we wish to avoid. None of the the results in this paper or in [16] see the difference between w and w^{-1} , so no changes in the statements of results need to be made to account for this difference.)

A finite Coxeter group (W, S) has a faithful action on $V(W) \cong \mathbb{R}^{|S|}$, known as the **reflection representation**, in which elements of S (and hence T) act as reflections. Absolute length has the following interpretation due to Carter [8, Lemma 2] in terms of the reflection representation.

Lemma 2.2. *Let $w \in W$, and let $A \subseteq V(W)$ be the subspace consisting of all points fixed by w . Then $\ell_T(w) = \text{codim } A = |S| - \dim A$.*

We will need the following simple corollary of this lemma.

Corollary 2.3. *Let $w \in W$, with $w = t_1 \cdots t_k$ for some $t_1, \dots, t_k \in T$. Suppose there exists a vector \mathbf{v} and an index i such that \mathbf{v} is fixed by w but not by t_i . Then $\ell_T(w) < k$.*

Proof. We prove the contrapositive. Suppose that $k = \ell_T(w)$, let A be the subspace fixed by w , and let H_{t_i} be the hyperplane fixed by t_i . Note $A \supseteq \bigcap_{i=1}^k H_{t_i}$. However, $\text{codim } A = k$, so $A = \bigcap_{i=1}^k H_{t_i}$, which means that every vector fixed by w must be fixed by every t_i . \square

Given $w \in W$, define the inversion arrangement

$$\mathcal{A}_w := \{H_t \subseteq V(W) \mid t \in \text{Inv}(w)\},$$

where

$$H_t := \{\mathbf{v} \in V(W) \mid t \cdot \mathbf{v} = \mathbf{v}\}$$

is the hyperplane fixed by t . As before, let $c(w)$ denote the number of chambers of \mathcal{A}_w . Hultman [16] extended Theorem 2.1 to an arbitrary finite Coxeter group using a condition on the **Bruhat graph** $\mathcal{B}(W)$ of W . This is the directed graph whose vertices correspond to the elements of W , with an edge from u to v if there exists $t \in T$ with $v = ut$ and $\ell(v) > \ell(u)$. (Note that v is not necessarily a cover of u in Bruhat order, since the difference in length may be greater than 1.)

There are two possible notions of distance in $\mathcal{B}(W)$. The **directed distance** $\ell_D(u, w)$ is the length of a shortest directed path from u to w in $\mathcal{B}(W)$, with $\ell_D(u, w) = \infty$ if there is no such path. The **undirected distance** $\ell_T(u, w)$ is the length of a shortest path from u to w ignoring the directions on the edges. Note that $\ell_T(u, w) \leq \ell_D(u, w)$ by definition, and a theorem of Dyer [10] states that $\ell_T(w) = \ell_T(\text{id}, w) = \ell_D(\text{id}, w)$ for all w . Furthermore, $\ell_T(u, w) = \ell_T(w^{-1}u, \text{id}) = \ell_T(w^{-1}u)$. Hultman's theorem [16] is the following.

Theorem 2.4. *Let (W, S) be a finite Coxeter group, and let $w \in W$. Then $c(w) = s(w)$ if and only if, for all $u \leq w$, $\ell_D(u, w) = \ell_T(u, w)$.*

Note that Hultman, Linusson, Shareshian, and Sjöstrand [15] had in fact shown that $c(w) \leq s(w)$ for arbitrary finite Coxeter groups. Hultman [16] goes on to give a much shorter and more conceptual proof of the difficult direction of Theorem 2.1(2) (that the right hull condition implies $c(w) = s(w)$) using Theorem 2.4. Our goal is to find the

analogues of the right hull condition and being defined by inclusions for B_n as well as a pattern avoidance criterion for the elements where equality holds. We will follow in outline the proof of Hultman.

Example 2.7. Let $w = 4231 \in S_4$, and let $u = 1324$. Note that $\ell_T(u, w) = 2$, since both u and w are adjacent to the identity in $\mathcal{B}(S_4)$. On the other hand, if $u < ut \leq w$, then $ut \in \{3124, 2314, 1423, 1342\}$, and no permutation in this set is adjacent to w in $\mathcal{B}(S_4)$, so $\ell_D(u, w) > 2$.

2.4. Billey–Postnikov avoidance. Billey and Postnikov [5] introduced a generalization of pattern avoidance particularly well suited to algebraic combinatorics on Coxeter groups. Let (W, S) be a finite Coxeter group. A **parabolic subgroup** is a subgroup $P \subseteq W$ that consists of all the elements fixing pointwise some subspace $A \subseteq V(W)$. In other words, P is parabolic if there exists $A \subseteq V(W)$, where $V(W)$ is the reflection representation, such that

$$P = \{w \in W \mid w \cdot \mathbf{v} = \mathbf{v} \text{ for all } \mathbf{v} \in A\}.$$

Let $\mathcal{B}(P)$ be the subgraph of the Bruhat graph $\mathcal{B}(W)$ induced by the vertices in P . Let $R \subseteq P$ be the elements corresponding to vertices with exactly one incoming edge in $\mathcal{B}(P)$. Then (P, R) is a Coxeter group, and we can define ℓ_P and \leq_P as length and Bruhat order in P (with respect to R). Note that $\ell_P(v) \leq \ell_W(v)$ for all $v \in P$ and, if $v \leq_P v'$, then $v \leq_W v'$. Since $v \leq_P v'$ implies $v \leq_W v'$, the Bruhat graph of P (with respect to R) is $\mathcal{B}(P)$ as the edges are correctly directed. Also, given $w \in P$, by Lemma 2.2, $\ell_T(w)$ is the same whether we consider w as an element of P or of W since, when considering w as an element of P , both $|S|$ and the codimension of the subspace fixed by w are decreased by $\dim A$. (The reader is cautioned that we have given a rather unusual definition of parabolic subgroups that has the drawback of applying only to the finite case. It can be seen to be equivalent to the usual definition by, for example, [18, Section 5.2]. We have taken more care than usual in selecting R as we need the positive roots of P to be the positive roots of W that are orthogonal to A .)

Example 2.8. Consider S_4 acting on \mathbb{R}^4 by permuting the coordinates. (Strictly speaking, the reflection representation of S_4 is the quotient of \mathbb{R}^4 by the line where $x_1 = x_2 = x_3 = x_4$, but this is irrelevant for us.) Consider the subspace A consisting of points such that $x_1 = x_2 = x_4$. This is fixed by the parabolic subgroup

$$P = \{1234, 2134, 1432, 2431, 4132, 4231\}$$

of elements w such that $w(3) = 3$, and $R = \{2134, 1432\}$.

Billey and Postnikov [5] define a **flattening map** $\text{fl}_P^W : W \rightarrow P$ as follows. Given $w \in W$, let $\text{Inv}(w)$ be the set of inversions of w . Then there is a unique element of P whose inversions are $\text{Inv}(w) \cap P$. We let $\text{fl}_P^W(w)$ be this element. Billey and Braden [4, Theorem 2] show that the flattening map satisfies the following. (To be precise, they state the theorem with multiplication backwards from what we have written here.)

Theorem 2.5. *Let W be a Coxeter group, $P \subseteq W$ a parabolic subgroup, and $\text{fl}_P^W : W \rightarrow P$ the flattening map. Then*

- (1) The map fl_P^W is P -equivariant, meaning that $\text{fl}_P^W(wv) = \text{fl}_P^W(w)v$ for all $v \in P$, $w \in W$.
- (2) If $\text{fl}_P^W(w) \leq_P \text{fl}_P^W(wv)$ for some $w \in W$, $v \in P$, then $w \leq_W wv$.

Now let (W', S') and (W, S) be arbitrary Coxeter groups, $v \in W'$, and $w \in W$. We say that w **BP contains** v if there exist a parabolic subgroup $P \subseteq W$ and an isomorphism $\phi : (W', S') \rightarrow (P, R)$ such that $\text{fl}_P^W(w) = \phi(v)$. Otherwise, w **BP avoids** v .

When $W = S_n$ and $W' = S_m$, the parabolic subgroups $P \subseteq W$ isomorphic to W' can all be constructed by selecting some indices $1 \leq i_1 < \dots < i_m \leq n$ and letting P be the subgroup

$$P = \{w \in S_n \mid w(j) = j \text{ for all } j \notin \{i_1, \dots, i_m\}\}.$$

In this case,

$$R = \{(i_j \ i_{j+1}) \mid 1 \leq j \leq m-1\}.$$

BP avoidance is almost the same as pattern avoidance for $W = S_n$. If we take the isomorphism $\phi : (S_m, S) \rightarrow (P, R)$ given by $\phi(s_j) = (i_j \ i_{j+1})$, then $v := \phi^{-1}(\text{fl}_P^{S_m}(w))$ is the permutation such that $v(j) > v(k)$ if and only if $w(i_j) > w(i_k)$, so w pattern contains v as usual. However, there is another Coxeter group isomorphism $\phi' : (S_m, S) \rightarrow (P, R)$ given by $\phi'(s_j) = (i_{m-j} \ i_{m-j+1})$ (reversing the Dynkin diagram). Since $\phi'(v) = \phi(w_0 v w_0)$, where $w_0 \in S_m$ is the element such that $w_0(i) = m+1-i$, we see that w BP contains v if and only if w pattern contains either v or $w_0 v w_0$.

Note that the converse of Theorem 2.5(2) holds for $W = S_n$ and $W' = S_m$, as can easily be seen from the tableau criterion.

Proposition 2.6. *Let $w \in S_n$, $v \in S_m$, $Q \subseteq S_n$ the subgroup of permutations which fix every entry other than i_1, \dots, i_m , and $\phi : S_m \rightarrow Q$ the order-preserving isomorphism. If $w \leq_{S_n} w\phi(v)$, then $\text{fl}_Q^{S_n}(w) \leq_Q \text{fl}_Q^{S_n}(w)\phi(v)$.*

2.5. Conventions for type B. We consider the group B_n as the following subgroup of the symmetric group S_{2n} , as in the book of Björner and Brenti [6, Section 8.1],

$$B_n := \{w \in S_{2n} \mid w(i) + w(2n+1-i) = 2n+1 \text{ for all } i, 1 \leq i \leq n\}.$$

We can equivalently restate this condition by saying that $w \in B_n$ if $w_0 w w_0 = w$.

The group B_n is a finite Coxeter group with simple generators $s_0 := (n \ n+1)$ and $s_i := (n-i \ n-i+1)(n+i \ n+i+1)$ for all i with $1 \leq i \leq n-1$. It acts on \mathbb{R}^n with s_0 acting by reflection across the hyperplane $x_1 = 0$ and s_i acting by reflection across the hyperplane $x_i - x_{i+1} = 0$.

Bruhat order on B_n turns out to be equal to the partial order induced from Bruhat order on S_{2n} under this embedding. (See, for example, [6, Cor. 8.1.9].) Given $w \in B_n$, we can define $E(w)$ by considering w as an element of S_{2n} . Because $w_0 w w_0 = w$, $E(w)$ has a rotational symmetry about $(n+1, n)$, so $(p, q) \in E(w)$ if and only if $(2n+2-p, 2n-q) \in E(w)$. Furthermore,

$$r_w(2n+2-p, 2n-q) = p - q - 1 + r_w(p, q),$$

which implies that, for $w \in B_n$, $r_w(p, q) = \max(0, q - p + 1)$ if and only if

$$r_w(2n + 2 - p, 2n - q) = \max(0, p - q - 1) = \max(0, (2n - q) - (2n + 2 - p) + 1).$$

We need slight weakenings of the right hull condition and the concept of being defined by inclusions. Given $w \in B_n$, we say that w is **defined by pseudo-inclusions** if, for all $(p, q) \in E(w)$, we have either $r_w(p, q) = \max(0, q - p + 1) = r_{\text{id}}(p, q)$ (as in the definition of being defined by inclusions) or $p = n + 1$, $q = n$, and $r_{n+1,n}(w) = 1$. Similarly, $w \in B_n$ satisfies the **relaxed right hull condition** if either, for every $u \in S_{2n}$ satisfying $u \subseteq H(w)$, we have $u \leq w$, or both $r_w(n + 1, n) = 1$ and, for every $u \in S_{2n}$ satisfying $u \subseteq H(w)$ and $r_u(n + 1, n) \leq 1$, we have $u \leq w$. (Note that it is not sufficient to consider only $u \in B_n$, as Example 4.1 shows. This is because the coessential set as we have defined it here is an S_{2n} concept and not truly appropriate for B_n , as discussed in Section 4.)

Example 2.9. Let $w = 362514 \in B_3$. Then $E(w) = \{(4, 1), (4, 3), (4, 5)\}$. We have $r_w(4, 1) = 0$, $r_w(4, 3) = 1$, and $r_w(4, 5) = 2$. The element w is not defined by inclusions since $r_w(4, 3) = 1$, but it is defined by pseudo-inclusions. Similarly, it does not satisfy the right hull condition but does satisfy the relaxed right hull condition.

The cycle decomposition of an element of B_n has a special structure. Since conjugation by w_0 fixes w for any $w \in B_n$, conjugating any cycle c in the cycle decomposition of w also gives a cycle $\bar{c} = w_0 c w_0$ of w . If $c = \bar{c}$, then we let $\mathbf{c} = c = \bar{c}$ and call \mathbf{c} an **odd cycle** of w . If $c \neq \bar{c}$, then we let $\mathbf{c} = c\bar{c}$ and call \mathbf{c} an **even cycle** of w . Alternatively, \mathbf{c} is an odd cycle if there is an odd number of i such that $i \leq n$ and $\mathbf{c}(i) > n$, and \mathbf{c} is an even cycle if there is an even number of i such that $i \leq n$ and $\mathbf{c}(i) > n$. It is easy to show that the reflection length of w (as an element of B_n) is $n - \text{ecyc}(w)$, where $\text{ecyc}(w)$ is the number of even cycles of w . (The cycle decomposition for elements of B_n and its relation to reflection length have appeared in the literature many times. Our description of the cycle decomposition comes from [3] and our terminology from [22]. The fact about reflection length is stated in [9] and [17]. Curiously, [9] uses the term “balanced” cycle to mean an even one, but [17] uses the term “balanced” cycle to mean an odd one!)

Billey–Postnikov avoidance in type B works as follows. There are two types of irreducible parabolic subgroups (by which we mean parabolic subgroups isomorphic to irreducible Coxeter groups) in B_n . Given indices $1 \leq i_1 < i_2 < \dots < i_m \leq 2n$ with the property that $i_j + i_k \neq 2n + 1$ for any j, k , we have an irreducible parabolic subgroup P isomorphic to S_m generated by $(i_j \ i_{j+1})(2n + 1 - i_{j+1} \ 2n + 1 - i_j)$ for all j with $1 \leq j \leq m - 1$. If our isomorphism $\phi : S_m \rightarrow P$ is given by

$$\phi(s_j) := (i_j \ i_{j+1})(2n + 1 - i_{j+1} \ 2n + 1 - i_j),$$

then $\text{fl}_P^{B_n}(w)$ is given by the relative order of $w(i_1), \dots, w(i_m)$. In particular, $\text{fl}_P^{B_n}(w) = \text{fl}_Q^{S_{2n}}(w)$, where Q is the parabolic subgroup of S_{2n} generated by the elements $(i_j \ i_{j+1})$. Hence, given $w \in B_n$, w BP contains $v \in S_m$ if w pattern contains v using some indices $i_1 < i_2 < \dots < i_m$ having the property that $i_j + i_k \neq 2n + 1$ for any j, k . (Note choosing indices $2n + 1 - i_m < \dots < 2n + 1 - i_1$ instead gets the same parabolic subgroup, but the isomorphism $\phi : S_m \rightarrow P$ is changed by conjugation with w_0 .)

Example 2.10. Let $w = 52863174 \in B_4$ and $v = 4231 \in S_4$. Taking $i_1 = 1$, $i_2 = 2$, $i_3 = 5$, and $i_4 = 6$, we see that w BP contains v .

Additionally, given indices $i_1 < \cdots < i_m < i_{m+1} < \cdots < i_{2m}$ with $i_j + i_{2m+1-j} = 2n + 1$ for all j , we have a parabolic subgroup P isomorphic to B_m generated by $(i_m \ i_{m+1})$ and $(i_j \ i_{j+1})(i_{2m-j} \ i_{2m+1-j})$ for all j with $1 \leq j \leq m - 1$. Our isomorphism $\phi : B_m \rightarrow P$ must be given by

$$\phi(s_0) := (i_m \ i_{m+1})$$

and

$$\phi(s_j) := (i_{m-j} \ i_{m-j+1})(i_{m+j} \ i_{m+j+1}),$$

and $\text{fl}_P^{B_n}(w)$ is given by the relative order of $w(i_1), \dots, w(i_{2m})$. Hence, $w \in B_n$ BP contains $v \in B_m$ if w pattern contains v using some indices $i_1 < \cdots < i_m < i_{m+1} < \cdots < i_{2m}$ with $i_j + i_{2m+1-j} = 2n + 1$ for all j .

Example 2.11. Let $w = 52863174 \in B_4$ and $v = 426153 \in B_3$. Taking $i_1 = 1$, $i_2 = 2$, $i_3 = 3$, $i_4 = 6$, $i_5 = 7$, and $i_6 = 8$, we see that w BP contains v .

Note that, given $w \in B_n$ and $v \in S_m$, whether w BP contains v or not depends on whether we consider v as an element of S_m or v as an element of B_m via, for example, the standard embedding of S_m in B_m . Similarly, if $v \in S_m$ with m even such that v happens to be an element of $B_{m/2}$, whether $w \in B_n$ BP contains v or not depends on whether we consider v as an element of S_m or as an element of $B_{m/2}$.

Example 2.12. Let $w = 52863174 \in B_4$. Then w BP contains $v = 4231 \in S_4$, but w does *not* BP contain $v = 4231 \in B_2$, nor does w BP contain $v = 42318675 \in B_4$.

Example 2.13. This example shows that Proposition 2.6 is false for BP avoidance in B_n , even when restricted to irreducible parabolic subgroups and even when $w \in Q$. Let $w = 426153 \in B_3$. Consider the parabolic Q generated by $R = \{r_1 = 132546, r_2 = 426153\}$, with the isomorphism $\phi : S_3 \rightarrow Q$ given by $\phi(s_1) = r_1$ and $\phi(s_2) = r_2$. (This is equivalent to taking $i_1 = 2$, $i_2 = 3$, and $i_3 = 6$.) Then $w = r_2$, and $u = r_1 = 132546 \leq_{B_3} w$, but $\phi^{-1}(u) = 213 \not\leq_{S_3} \phi^{-1}(w) = 132$.

3. PROOF OF MAIN THEOREM

We now state our theorem and outline the proof, deferring individual details to subsections.

Main Theorem. *Let $w \in B_n$. Then the following are equivalent.*

- (1) *The number of chambers of the inversion arrangement \mathcal{A}_w is equal to the number of elements in $[\text{id}, w]$.*
- (2) *For any $u \leq w$, the directed distance from u to w in the Bruhat graph is the same as the undirected distance from u to w .*
- (3) *The element w is defined by pseudo-inclusions.*
- (4) *The element w satisfies the relaxed right hull condition.*

- (5) *The element w BP avoids* $4231 \in S_4$, 35142 , $42513 \in S_5$, $351624 \in S_6$, 563412 , 653421 , 645231 , 635241 , 624351 , 642531 , 536142 , 426153 , 462513 , $623451 \in B_3$, 47618325 , 46718235 , 57163824 , 37581426 , 47163825 , 46172835 , 37518426 , 35718246 , 37145826 , 37154826 , 52618374 , 42681375 , 42618375 , $35172846 \in B_4$, and $3517294a68$, $3517924a68$, $3617294a58 \in B_5$.

We use the notation $a = 10$ in elements of B_5 to avoid confusion. In what follows, we will call elements that satisfy Condition 2 **Hultman elements**.

Proof. Conditions 1 and 2 are equivalent by Theorem 2.4.

Corollary 3.4 shows that Conditions 3 and 4 are equivalent.

Suppose Condition 4 holds, so $w \in B_n$ satisfies the relaxed right hull criterion. Then, by Proposition 3.5, w is Hultman.

Suppose Condition 3 does not hold, so $w \in B_n$ is not defined by pseudo-inclusions. Then, by Proposition 3.6, w BP contains some $v \in B_m$ or S_{m+1} , with $m \leq 5$, such that v is not defined by pseudo-inclusions. By computer calculation, v BP contains one of the elements listed in Condition 5. Since BP containment is transitive, Condition 5 does not hold for w either.

None of the patterns listed in Condition 5 are Hultman, as shown by the data in Figure 3. Hence, by Proposition 3.7, if $w \in B_n$ does not satisfy Condition 5, w is not Hultman. \square

3.1. The right hull condition and being defined by inclusions. In this section, we show that Sjöstrand’s right hull condition and the Gasharov–Reiner condition of being defined by inclusions are equivalent for permutations. We first prove the following lemma, which is a restatement of [26, Lemma 3.2].

Lemma 3.1. *Let $w \in S_n$ and $1 \leq p, q \leq n$. Then the following are equivalent:*

- (1) $r_w(p, q) = q - p + 1$.
- (2) $w(k) \geq p$ for all $k > q$.
- (3) $\{1, \dots, p - 1\} \subseteq \{w(1), \dots, w(q)\}$.

Proof. Since w is a bijection considered as a function $w : \{1, \dots, n\} \rightarrow \{1, \dots, n\}$, the second and third conditions are equivalent. Now note that

$$\begin{aligned} n - p + 1 &= \#\{k \mid w(k) \geq p\} \\ &= \#\{k \leq q \mid w(k) \geq p\} + \#\{k > q \mid w(k) \geq p\} \\ &= r_w(p, q) + \#\{k > q \mid w(k) \geq p\}, \end{aligned}$$

so $r_w(p, q) = q - p + 1$ if and only if $\#\{k > q \mid w(k) \geq p\} = n - q$, and the last statement holds if and only if $w(k) \geq p$ for all $k > q$. \square

To show the equivalence of the right hull condition and being defined by inclusions, we prove the following slightly stronger statement.

Proposition 3.2. *Let $w \in S_n$. Then $u \subseteq H(w)$ if and only if, for all $(p, q) \in E(w)$ with $r_w(p, q) = \max(0, q - p + 1)$, $r_u(p, q) \leq r_w(p, q)$.*

Proof. First we assume that $u \not\subseteq H(w)$ and show that $r_u(p, q) > r_w(p, q)$ for some $(p, q) \in E(w)$ with $r_w(p, q) = \max(0, q - p + 1)$. If $u \not\subseteq H(w)$, either there exists some i such that $w(j) < u(i)$ or $j > i$ for all j , or there exists some i such that $w(j) > u(i)$ or $j < i$ for all j , or both. In the first case, let j_1 be the largest j such that $j_1 \leq i$ and no k satisfies both $k < j_1$ and $w(k) > w(j_1)$. Note that j_1 exists since $j = 1$ satisfies both conditions. Furthermore, let j_2 be the smallest j such that $j_2 > i$ and no k satisfies both $k < j_2$ and $w(k) > w(j_2)$. If $w^{-1}(n) \leq i$, then neither $n < u(i)$ nor $w^{-1}(n) > i$ would hold, so $j = w^{-1}(n)$ satisfies both conditions, and j_2 exists. Then $(w(j_1) + 1, j_2 - 1) \in E(w)$, and $r_w(w(j_1) + 1, j_2 - 1) = 0$, but $r_u(w(j_1) + 1, j_2 - 1) \neq 0$ since $i \leq j_2 - 1$ and $u(i) \geq w(j_1) + 1$.

In the second case, let j_1 be the largest j such that $j_1 < i$ and no k satisfies both $k > j_1$ and $w(k) < w(j_1)$. Here j_1 exists since $j = w^{-1}(1)$ satisfies both conditions. Also, let j_2 be the smallest j such that $j_2 \geq i$ and no k satisfies both $k > j_2$ and $w(k) < w(j_2)$; $j = n$ satisfies both conditions, so j_2 exists. Then $(w(j_2), j_1) \in E(w)$. Furthermore, there is no k with $w(k) < w(j_2)$ and $k > j_1$, which, by Lemma 3.1, implies $r_w(w(j_2), j_1) = j_1 - w(j_2) + 1$. Also by Lemma 3.1, $r_u(w(j_2), j_1) > r_w(w(j_2), j_1)$.

Now we assume that $u \subseteq H(w)$ and show that $r_u(p, q) = r_w(p, q)$ for all (p, q) such that $r_w(p, q) = \max(0, q - p + 1)$. (We do not need to assume for this part that $(p, q) \in E(w)$, so in fact we prove a stronger statement.) Suppose $r_w(p, q) = 0$. Then $w(k) < p$ for all $k \leq q$. Hence, if $u \subseteq H(w)$, $u(i) < p$ for all $i \leq q$ since, otherwise, there would exist $k' \leq k \leq q$ with $w(k') \geq u(k) \geq p$, contradicting our assumption that $r_w(p, q) = 0$. Therefore, $r_u(p, q) = 0$.

Similarly, suppose $r_w(p, q) = q - p + 1$. By Lemma 3.1, $w(k) \geq p$ for all $k > q$. Hence, if $u \subseteq H(w)$, we also have $u(k) \geq p$ for all $k > q$ since, otherwise, there would exist $k' \geq k > q$ with $w(k') \leq u(k) < p$. Therefore, $r_u(p, q) = q - p + 1$. \square

Corollary 3.3. *A permutation w satisfies the right hull condition if and only if w is defined by inclusions.*

Proof. A permutation w fails to be defined by inclusions if and only if there exists $(a, b) \in E(w)$ such that $r_w(a, b) \neq \max(0, b - a + 1)$. Since $E(w)$ is the unique minimal set determining $[\text{id}, w]$, this means that w fails to be defined by inclusions if and only if there exists $u \not\subseteq w$ with $r_u(p, q) \leq r_w(p, q)$ for all $(p, q) \in E(w)$ such that $r_w(p, q) = \max(0, q - p + 1)$. (At least some such u will satisfy $r_u(a, b) > r_w(a, b)$.) By Proposition 3.2, $u \subseteq H(w)$, so w fails to be defined by inclusions if and only if w fails to satisfy the right hull condition. \square

Corollary 3.4. *Let $w \in S_{2n}$. Then w satisfies the relaxed right hull condition if and only if w is defined by pseudo-inclusions.*

Proof. A permutation $w \in S_{2n}$ fails to be defined by pseudo-inclusions if and only if there exists $(a, b) \in E(w)$ such that $r_w(a, b) \neq \max(0, b - a + 1)$ and $(a, b, r_w(a, b)) \neq (n + 1, n, 1)$. Since $E(w)$ is the unique minimal set determining $[\text{id}, w]$, this means that w fails to be defined by pseudo-inclusions if and only if there exists $u \not\subseteq w$ satisfying both the condition

that $r_u(p, q) \leq r_w(p, q)$ for all $(p, q) \in E(w)$ such that $r_w(p, q) = \max(0, q - b + 1)$ and, if $r_w(n + 1, n) = 1$, the condition that $r_u(n + 1, n) \leq 1$. By Proposition 3.2, $u \subseteq H(w)$, and, if $r_w(n + 1, n) = 1$, $r_u(n + 1, n) \leq 1$ also, so, by definition, w fails to satisfy the relaxed right hull condition. \square

3.2. Defined by pseudo-inclusion elements are Hultman. In this section, we prove the following proposition.

Proposition 3.5. *Suppose $w \in B_n$ satisfies the relaxed right hull condition. Then $\ell_D(u, w) = \ell_T(u, w)$ for all $u \leq w$.*

Our proof largely follows that of Hultman [16, Lemma 4.4] showing that permutations satisfying the right hull condition are Hultman, but it is significantly more complicated because Proposition 2.6 does not hold for B_n and arguments to work around this issue are necessary.

Proof. Let $w \in B_n$ satisfy the relaxed right hull condition. We need to show that, given any $u \leq w$, we have $\ell_D(u, w) = \ell_T(u, w)$. By induction on $\ell(w) - \ell(u)$, it suffices to show that there exists a reflection t with $u < ut \leq w$ and $\ell_T(ut, w) = \ell_T(w^{-1}ut, \text{id}) = \ell_T(w^{-1}u, \text{id}) - 1$.

We call a cycle **trivial** if it consists simply of a pair of fixed points and **nontrivial** otherwise. First suppose that $w^{-1}u$ has no nontrivial even cycles and $u < w$. Let P be the parabolic subgroup generated by the reflections $(i \ 2n+1-i)$ and $(i \ j)(2n+1-i \ 2n+1-j)$ where $w^{-1}u(i) \neq i$ and $w^{-1}u(j) \neq j$, or equivalently $u(i) \neq w(i)$ and $u(j) \neq w(j)$. Observe that $uP = wP$. Let $\tilde{u} = \text{fl}_P^{B_n}(u)$ and $\tilde{w} = \text{fl}_P^{B_n}(w)$. Since, in this case, $\text{fl}_P^{B_n}(w) = \text{fl}_Q^{S_{2n}}(w)$ where Q is the parabolic subgroup of S_{2n} generated by the reflections $(i \ j)$ for which $u(i) \neq w(i)$ and $u(j) \neq w(j)$, we have $\tilde{u} < \tilde{w}$ by Proposition 2.6 (and the fact that Bruhat order in B_n is induced from Bruhat order for S_{2n}). Hence, by the definition of Bruhat order, there exists a reflection $t \in P$ such that $\tilde{u} <_P \tilde{u}t \leq_P \tilde{w}$. By Theorem 2.5, $u < ut \leq w$. Since all nontrivial cycles of $w^{-1}u$ are odd, $w^{-1}u$ has maximal reflection length in P , so $\ell_T(w^{-1}ut, \text{id}) = \ell_T(w^{-1}u, \text{id}) - 1$.

Now suppose there is a nontrivial even cycle $\mathbf{c} = c\bar{c}$ in the cycle decomposition of $w^{-1}u$. We first show $wc < w$ in Bruhat order on S_{2n} . (Note $wc \notin B_n$.) First note that $wc(i) = u(i)$ or $wc(i) = w(i)$ for all i . Since $u < w$, $(u(i), i) \in H(w)$ for all i , so $(wc(i), i) \in H(w)$ for all i . Hence, if w satisfies the (unrelaxed) right hull condition, $wc < w$. Otherwise, $r_w(n + 1, n) = 1$ and $r_u(n + 1, n) \leq 1$. If $r_u(n + 1, n) = 0$, then $r_{wc}(n + 1, n) \leq 1$. If $r_u(n + 1, n) = 1$, then let a be the unique index such that $a \leq n$ and $w(a) > n$, with $\alpha := w(a)$, and let b be the unique index such that $b \leq n$ and $u(b) > n$, with $\beta := u(b)$. Now, if $a = b$ or $\alpha = \beta$, then $r_{wc}(n + 1, n) = 1$, so $wc < w$. Otherwise, if $c(b) = b$, then $(wc)(b) = w(b) \leq n$, so $r_{wc}(n + 1, n) \leq 1$.

Finally, we are left with the case where $r_u(n + 1, n) = r_w(n + 1, n) = 1$, $a \neq b$, $\alpha \neq \beta$, and $c(b) \neq b$. Since $a \neq b$, $c(b) = w^{-1}(\beta) > n$. We now show that $c(a) \neq a$. Note that, as $b \leq n$, there must exist some $b' > n$ such that $c(b') \leq n$. Furthermore, $b' \neq 2n + 1 - b$, since \mathbf{c} is even, so $c(2n + 1 - b) = 2n + 1 - b$. Since $r_u(n + 1, n) = 1$ and $u \in B_n$, we can only have $i > n$ and $u(i) \leq n$ if $i = 2n + 1 - b$, so $u(b') > n$. It follows that, as

$c(b') = w^{-1}(u(b')) \leq n$, $u(b') = \alpha$ and $c(b') = a$, so $c(a) \neq a$. Hence, $wc(a) = u(a)$, and the only index $i \leq n$ with $wc(i) > n$ is $i = b$. Therefore, $r_{wc}(n+1, n) = 1$. Since we also have $wc \in H(w)$, and w satisfies the relaxed right hull property, $wc < w$.

Now let $i_1 < i_2 \cdots < i_m$ be the indices such that $c(i_j) \neq i_j$. (In particular, we let m be the number of such indices.) Consider the parabolic subgroup $P_c \subseteq B_n$ with the isomorphism $\phi : S_m \rightarrow P_c$ given by $\phi(s_j) = (i_j \ i_{j+1})(2n+1-i_{j+1} \ 2n+1-i_j)$. Note $\mathbf{c} \in P_c$ and, indeed, \mathbf{c} is an element of maximal reflection length in P_c , so for any reflection $t \in P_c$, $\ell_T(w^{-1}ut, \text{id}) = \ell_T(w^{-1}u, \text{id}) - 1$.

Moreover, $\text{fl}_{P_c}^{B_n}(w\mathbf{c}) = \text{fl}_Q^{S_{2n}}(wc)$, and $\text{fl}_{P_c}^{B_n}(w) = \text{fl}_Q^{S_{2n}}(w)$, where Q is the parabolic subgroup of S_{2n} generated by the elements $(i_j \ i_{j+1})$. Let $\tilde{u} = \text{fl}_{P_c}^{B_n}(u)$ and $\tilde{w} = \text{fl}_{P_c}^{B_n}(w)$. As $wc < w$, by Theorem 2.6, $\text{fl}_Q^{S_{2n}}(wc) < \text{fl}_Q^{S_{2n}}(w)$, and

$$\tilde{u} = \text{fl}_{P_c}^{B_n}(u) = \text{fl}_{P_c}^{B_n}(w\mathbf{c}) = \text{fl}_Q^{S_{2n}}(wc) < \text{fl}_Q^{S_{2n}}(w) = \text{fl}_{P_c}^{B_n}(w) = \tilde{w}.$$

By the definition of Bruhat order, there exists $t \in P_c$ such that $\tilde{u} <_{P_c} \tilde{u}t \leq_{P_c} \tilde{w}$. By Proposition 2.5, $u < ut$.

We now show that $ut \leq w$. For i such that $\mathbf{c}(i) \neq i$, since $\tilde{u}t \leq \tilde{w}$,

$$(i, ut(i)) \in H(\{(w(i), i) \mid \mathbf{c}(i) \neq i\}) \subseteq H(w).$$

For i such that $\mathbf{c}(i) = i$, we know $ut(i) = u(i)$, so $(ut(i), i) \in H(w)$ since $u \leq w$. Hence, in the case where w satisfies the (unrelaxed) right hull condition, $ut \leq w$.

Otherwise, $r_w(n+1, n) = 1$, and $r_u(n+1, n) \leq 1$. Let $t = (j \ k)(2n+1-j \ 2n+1-k) \in P_c$, and assume without loss of generality that $j < k$ and that $c(j) \neq j$ while $\bar{c}(j) = j$. Assume for contradiction that $r_{ut}(n+1, n) > r_u(n+1, n)$. Then $j, u(j) \leq n$, and $k, u(k) > n$. (We would also have $r_{ut}(n+1, n) > r_u(n+1, n)$ if $2n+1-k, u(2n+1-k) \leq n$ and $2n+1-j, u(2n+1-j) > n$, but that amounts to the same condition.) But then, if $j, u(j) \leq n$ and $k, u(k) > n$, we would have $k > n$ and $ut(k) \leq n$, while $j \leq n$ and $ut(j) > n$. However, this implies

$$\{(ut(j), j), (ut(k), k)\} \subseteq H(\{(w(i), i) \mid c(i) \neq i\})$$

since $\tilde{u}t \leq_{P_c} \tilde{w}$. As before, let a be the unique index such that $a \leq n$ and $w(a) > n$, which implies that $2n+1-a$ is the unique index with $2n+1-a > n$ and $w(2n+1-a) \leq n$. Since \mathbf{c} is an even cycle, either $c(a) = a$ or $c(2n+1-a) = 2n+1-a$. In the first case, there does not exist any i with $c(i) \neq i$ such that $i \leq n$ and $w(i) > n$, so

$$(ut(j), j) \notin H(\{(w(i), i) \mid c(i) \neq i\}),$$

a contradiction, and, in the second case,

$$(ut(k), k) \notin H(\{(w(i), i) \mid c(i) \neq i\}).$$

Therefore, $r_{ut}(n+1, n) = r_u(n+1, n) \leq r_w(n+1, n)$. Since w satisfies the relaxed right hull condition, $ut \leq w$. \square

3.3. Failure of inclusion conditions and pattern containment. Now we show that an element that fails to be defined by pseudo-inclusions must not be Hultman. We first show that any element not defined by pseudo-inclusions BP contains one of 27 elements. We give a short proof here that relies on a computer calculation. A longer proof by hand involving a tedious case-by-case analysis of the possible ways we can have $i_j + i_k = 2n + 1$ in the notation of the following proof is possible.

Proposition 3.6. *Suppose $w \in B_n$ is not defined by pseudo-inclusions. Then w BP contains some $v \in W$, where W is either S_{m+1} or B_m for some $m \leq 5$, such that v is not defined by pseudo-inclusions.*

Proof. If $w \in B_n$ is not defined by pseudo-inclusions, then either $(n + 1, n) \in E(w)$ and $r_w(n + 1, n) \geq 2$, or $(p, q) \in E(w)$ (with $(p, q) \neq (n + 1, n)$) and $r_w(p, q) \geq \max(1, q - p + 2)$.

Suppose $(n + 1, n) \in E(w)$ and $r_w(n + 1, n) \geq 2$. Since $r_w(n + 1, n) \geq 2$, there exist $i_1, i_2 \leq n$ with $w(i_1), w(i_2) > n$. Let $i_3 = w^{-1}(n)$ and $i_4 = n$. Since $(n + 1, n) \in E(w)$, $i_3, w(i_4) \leq n$. Let P be the parabolic subgroup generated by the (not necessarily simple) reflections $(i_j \ i_k)(2n + 1 - i_k \ 2n + 1 - i_j)$ and $(i_4 \ 2n + 1 - i_4)$. Depending on whether $i_3 = i_4$ or not, $P \cong B_3$ or $P \cong B_4$; let $m = 3$ or $m = 4$ respectively, and let $\phi : B_m \rightarrow P$ be the isomorphism. Consider $v = \phi^{-1}(\text{fl}_P^{B_n}(w))$. Since $i_3 < n$ and $w(i_3) = n$, we have $v^{-1}(m) \leq m$, and since $i_4 = n$ with $w(i_4) < n$, we have $v(m) \leq m$. This implies that $(m + 1, m) \in E(v)$. Furthermore, $r_v(m + 1, m) \geq 2$ since (the flattenings of) the points at i_1 and i_2 both count towards $r_v(m + 1, m)$.

Now suppose $(p, q) \in E(w)$ with $(p, q) \neq (n + 1, n)$ and $r_w(p, q) \geq \max(1, q - p + 2)$. Since $r_w(p, q) \geq \max(1, q - p + 2)$, we must have some $i_1 \leq q$ with $w(i_1) \geq p$ and some $i_2 > q$ with $w(i_2) < p$. Let $i_3 = w^{-1}(p - 1)$, $i_6 = w^{-1}(p)$, $i_4 = q$, and $i_5 = q + 1$. Since $(p, q) \in E(w)$, $i_3 \leq i_4 < i_5 \leq i_6$, and $w(i_4) \leq w(i_3) < w(i_6) \leq w(i_5)$.

First we consider the case where $i_j + i_k \neq 2n + 1$ for any j, k with $1 \leq j, k \leq 6$. Let P be the parabolic subgroup generated by the (not necessarily simple) reflections $(i_j \ i_k)(2n + 1 - i_k \ 2n + 1 - i_j)$. Depending on whether $i_3 = i_4$ and $i_5 = i_6$, $P \cong S_{m+1}$ for some m , $3 \leq m \leq 5$, with some isomorphism $\phi : S_{m+1} \rightarrow P$. Consider $v = \phi^{-1}(\text{fl}_P^{B_n}(w))$. Note (p, q) will correspond to a box $(\tilde{p}, \tilde{q}) \in E(v)$, and $(w(i_1), i_1)$ and $(w(i_2), i_2)$ will force $r_v(\tilde{p}, \tilde{q}) \geq \max(1, \tilde{q} - \tilde{p} + 2)$, so v is not defined by inclusions.

If $i_j + i_k = 2n + 1$ for some j, k , then let P be the parabolic subgroup generated by $(i_j \ i_k)(2n + 1 - i_k \ 2n + 1 - i_j)$ and $(i_j \ 2n + 1 - i_j)$. Depending on how many coincidences of indices we have, $P \cong B_m$ for some m , $3 \leq m \leq 5$. (We cannot have $m = 2$, because if $i_3 = i_4 = 2n + 1 - i_5 = 2n + 1 - i_6$, then $(p, q) = (n + 1, n)$.) Again (p, q) will correspond to a box $(\tilde{p}, \tilde{q}) \in E(v)$. We cannot have $(\tilde{p}, \tilde{q}) = (m + 1, m)$ because, in that case, we would have had $w(i_3) \leq n < w(i_6)$ and $i_4 \leq n < i_5$, which would imply $(p, q) = (n + 1, n)$. Also, $(w(i_1), i_1)$ and $(w(i_2), i_2)$ will force $r_v(\tilde{p}, \tilde{q}) \geq \max(1, \tilde{q} - \tilde{p} + 2)$.

Hence, in all cases, w BP contains some v that is not defined by pseudo-inclusions. \square

Using a computer program, we find all the elements of S_{m+1} and B_m with $m \leq 5$ that are not defined by (pseudo)-inclusions. We then find among these elements the ones that

do not BP contain some other element not defined by (pseudo)-inclusions. These elements w are listed in the table in Figure 3.

For each w in Figure 3, we list all the elements $u < w$ such that $\ell_D(u, w) > \ell_T(u, w)$. Note there is at least one such u for each w , so none of these elements w are Hultman.

3.4. Failure of containment and Hultman's condition. To complete our proof, we show the following.

Proposition 3.7. *Suppose $w \in W$ BP contains $v \in W'$ and v is not Hultman. Then w is not Hultman.*

Proof. By definition, we have some parabolic subgroup $P \subseteq W$ with an isomorphism $\phi : W' \rightarrow P$ such that $\text{fl}_P^W(w) = \phi(v)$. Since v is not Hultman, there exists $u \leq_{W'} v$ such that $\ell_D(u, v) > \ell_T(u, v)$. Now let $x = w\phi(v^{-1}u)$. Note that

$$\ell_T(x, w) = \ell_T(w^{-1}x, \text{id}) = \ell_T(w^{-1}x) = \ell_T(\phi(v^{-1}u)) = \ell_T(v^{-1}u) = \ell_T(u, v).$$

Now consider a directed path

$$x = x_0, x_1, \dots, x_k = w$$

of length k in $\mathcal{B}(W)$. If $x_j \in wP$ for all j , then $\text{fl}_P^W(x_i) = \text{fl}_P^W(w)(w^{-1}x_i) = \phi(v)w^{-1}x_i$ for all i , and we have a directed path

$$u = \phi^{-1}(\text{fl}_P^W(x_0)), \phi^{-1}(\text{fl}_P^W(x_1)), \dots, \phi^{-1}(\text{fl}_P^W(x_k)) = v$$

from u to v in $\mathcal{B}(W')$. This is a path in $\mathcal{B}(W')$ because, for all i , $x_i = x_{i-1}t_i$ for some $t_i \in T \cap P$, so $v\phi^{-1}(w^{-1}x_i) = v\phi^{-1}(w^{-1}x_{i-1})\phi^{-1}(t_i)$. This path is appropriately directed, meaning $v\phi^{-1}(w^{-1}x_i) >_{W'} v\phi^{-1}(w^{-1}x_{i-1})$ as, otherwise, we would have $\text{fl}_P^W(x_i) <_P \text{fl}_P^W(x_{i-1})$, and, hence by Theorem 2.5, since $x_i = x_{i-1}t_i$ and $t_i \in P$, we would have $x_i < x_{i-1}$, which is false by assumption. Therefore, since k is the length of a directed path from u to v in $\mathcal{B}(W')$, $k \geq \ell_D(u, v) > \ell_T(u, v) = \ell_T(x, w)$.

On the other hand, suppose $x_j \notin wP$ for some j . For convenience, we choose the first such j , so $x_{j'} \in wP$ for all $j' < j$. Let $t_i = x_i^{-1}x_{i-1}$ for all i , $1 \leq i \leq k$. Then $t_j \notin P$. However, $w^{-1}x = t_k \cdots t_1 \in P$. Hence, by Corollary 2.3, $\ell_T(w^{-1}x) < k$.

Since any directed path from x to w in $\mathcal{B}(W)$ has length greater than $\ell_T(x, w)$, we must have that w is not Hultman. \square

4. COESSENTIAL SETS

The purpose of this section is to describe the relation between our results and the Coxeter-theoretic coessential set defined in [23].

Let W be an arbitrary Coxeter group, and let $w \in W$. The **(Coxeter-theoretic) coessential set** of w , denoted $\mathcal{E}(w)$, is the set of minimal elements of

$$\{v \in W \mid v \not\leq w\}.$$

| W | w | u | $\ell_D(u, w)$ | $\ell_T(u, w)$ |
|-------|------------|------------|----------------|----------------|
| S_4 | 4231 | 2143 | 4 | 2 |
| S_5 | 35142 | 12435 | 5 | 3 |
| S_5 | 42513 | 13245 | 5 | 3 |
| S_6 | 351624 | 423156 | 6 | 4 |
| S_6 | 351624 | 126543 | 6 | 4 |
| B_3 | 426153 | 132546 | 4 | 2 |
| B_3 | 536142 | 142536 | 4 | 2 |
| B_3 | 563412 | 124356 | 4 | 2 |
| B_3 | 462513 | 135246 | 4 | 2 |
| B_3 | 635241 | 153426 | 4 | 2 |
| B_3 | 635241 | 241635 | 4 | 2 |
| B_3 | 642531 | 315264 | 4 | 2 |
| B_3 | 642531 | 153426 | 4 | 2 |
| B_3 | 645231 | 154326 | 4 | 2 |
| B_3 | 645231 | 351624 | 4 | 2 |
| B_3 | 623451 | 132546 | 4 | 2 |
| B_3 | 624351 | 135246 | 4 | 2 |
| B_3 | 624351 | 142536 | 4 | 2 |
| B_3 | 653421 | 214365 | 4 | 2 |
| B_4 | 35172846 | 12436578 | 5 | 3 |
| B_4 | 46172835 | 12536478 | 5 | 3 |
| B_4 | 57163824 | 14627358 | 5 | 3 |
| B_4 | 57163824 | 12654378 | 5 | 3 |
| B_4 | 47163825 | 13527468 | 5 | 3 |
| B_4 | 47163825 | 12645378 | 5 | 3 |
| B_4 | 52618374 | 14236758 | 5 | 3 |
| B_4 | 52618374 | 13254768 | 5 | 3 |
| B_4 | 47618325 | 13254768 | 5 | 3 |
| B_4 | 42681375 | 13427568 | 5 | 3 |
| B_4 | 42681375 | 13254768 | 5 | 3 |
| B_4 | 42618375 | 13245768 | 5 | 3 |
| B_4 | 37154826 | 12536478 | 5 | 3 |
| B_4 | 37154826 | 12463578 | 5 | 3 |
| B_4 | 37145826 | 12436578 | 5 | 3 |
| B_4 | 37581426 | 14627358 | 5 | 3 |
| B_4 | 37581426 | 12654378 | 5 | 3 |
| B_4 | 37518426 | 12645378 | 5 | 3 |
| B_4 | 37518426 | 14263758 | 5 | 3 |
| B_4 | 35718246 | 12463578 | 5 | 3 |
| B_4 | 46718235 | 12354678 | 5 | 3 |
| B_5 | 3617294a58 | 124365879a | 6 | 4 |
| B_5 | 3617294a58 | 125347869a | 6 | 4 |
| B_5 | 3517924a68 | 124365879a | 6 | 4 |
| B_5 | 3517924a68 | 124538679a | 6 | 4 |
| B_5 | 3517294a68 | 124356879a | 6 | 4 |

FIGURE 3. BP-containment-minimal non-Hultman elements in types A and B

An element of W is **basic** if it is an element of $\mathcal{E}(w)$ for some $w \in W$. Hence, the set $\mathbf{B}(W)$ of all basic elements of W is

$$\mathbf{B}(W) := \bigcup_{w \in W} \mathcal{E}(w).$$

An element $u \in W$ is **bigrassmannian** if all reduced expressions for u share the same initial and final element. In other words, given $u = s_1 \cdots s_{\ell(u)} = s'_1 \cdots s'_{\ell(u)}$, then $s_1 = s'_1$ and $s_{\ell(u)} = s'_{\ell(u)}$. It is a theorem of Lascoux and Schützenberger [19] (with a subsequent different proof due to Geck and Kim [13]) that basic elements are bigrassmannian. The converse is true for $W = S_n$ but not in general.

For $W = S_n$, elements of $\mathcal{E}(w)$ correspond to elements of $E(w)$ as follows. Given a box at (p, q) with $r = r_w(p, q)$, we have a unique minimal permutation $v = v(p, q, r)$ with $r_v(p, q) = r + 1$. To be precise,

$$v(p, q, r) = 1 \cdots (q - r - 1)p \cdots (p + r)(q - r) \cdots (p - 1)(p + r + 1) \cdots n,$$

written in 1-line notation. The following proposition relates properties of the Coxeter-theoretic coessential set of an element $w \in S_n$ with the property of w being defined by inclusions.

Proposition 4.1. *Let $w \in S_n$. An element $(p, q) \in E(w)$ corresponds to an element of $\mathcal{E}(w)$ with a unique reduced expression if and only if $r_w(p, q) = r_{\text{id}}(p, q) = \max(0, q - p + 1)$.*

Proof. The element

$$v(p, q, r) = 1 \cdots (q - r - 1)p \cdots (p + r)(q - r) \cdots (p - 1)(p + r + 1) \cdots n$$

has a unique reduced expression precisely if $p = p + r$, in which case $r = 0$ and

$$v(p, q, r) = s_{p-1} \cdots s_{q-r} = s_{p-1} \cdots s_q$$

as a product of consecutive (in the Dynkin diagram) simple reflections, or if $p - 1 = q - r$, in which case $r = q - p + 1$ and

$$v(p, q, r) = s_{q-r} \cdots s_{p+r-1} = s_{p-1} \cdots s_q.$$

The two cases differ by whether the indices are increasing or decreasing. (Note that, if $q - p + 1 > 0$, so $q \geq p$, then $r = 0$ is not possible.) \square

This means that w is defined by inclusions if and only if every element of $\mathcal{E}(w)$ has a unique reduced expression.

Given $W = B_n \subseteq S_{2n}$ and $w \in W$, the elements of $E(w)$ (considering w as an element of S_{2n}) come in pairs that are rotationally symmetric around $(n + 1, n)$. In particular, if $(p, q) \in E(w)$, then $(2n + 2 - p, 2n - q) \in E(w)$, with

$$r_w(2n + 2 - p, 2n - q) = q - p + 1 + r_w(p, q).$$

However, in some cases, the set $E(w)$ may not be minimal for the purposes of determining if $v \leq w$ for $v \in B_n$. In particular, a pair

$$\{(p, q), (2n + 2 - p, 2n - q)\} \subseteq E(w)$$

| | $v(p, q, r)$ |
|-------------------------|---|
| $p + r \leq n$ | $1 \cdots (q - r - 1)p \cdots (p + r)(q - r) \cdots (p - 1)(p + r + 1) \cdots n$ |
| $p \leq n < p + r$ | $1 \cdots (q - r - 1)p \cdots n(2n + 2 - p) \cdots (n + r + 1)(q - r) \cdots (n - r - 1)$ |
| $n < p, p + q < 2n + 2$ | $1 \cdots (q - r - 1)p \cdots (p + r)(q - r) \cdots (2n - p - r)(2n + 2 - p) \cdots n$ |
| $p + q \geq 2n + 2$ | $1 \cdots (2n - p - r)(2n + 2 - p) \cdots qp \cdots (p + r)(q + 1) \cdots n$ |

FIGURE 4. B_n elements corresponding to coessential boxes where $q < n$ or both $q = n$ and $p > n$.

may not be needed to determine if $v \leq w$. For any $v \in B_n$, if $r_v(2n + 2 - p, q) \leq r$ with $p, q \leq n$, then, since either $v(i) > n \geq q$ or $v(2n + 1 - i) > n \geq q$ for any i , and in particular for i with $p \leq i \leq n$ (so $n < 2n + 1 - i < 2n + 2 - p$), we must have $r_v(p, q) \leq r - (n - p + 1)$. Suppose for some $p, q \leq n$, both $(p, q) \in E(w)$ and $(2n + 2 - p, q) \in E(w)$, and $r_w(2n + 2 - p, q) = r_w(p, q) + p - q - 1$. Then, in this case, for any $v \in B_n$, if $r_v(2n + 2 - p, q) \leq r_w(2n + 2 - p, q)$, then automatically $r_v(p, q) \leq r_w(p, q)$. Hence, to check if $v \leq w$, it is not necessary to explicitly check if $r_v(p, q) \leq r_w(p, q)$.

Given $w \in B_n$, let $E'(w) = E(w) \setminus S$, where S is the set of all redundant essential boxes. To be precise, S contains all boxes $(p, q), (2n + 2 - p, 2n - q) \in E(w)$ where $p, q < n$, $(2n + 2 - p, q) \in E(w)$, and $r_w(2n + 2 - p, q) = r_w(p, q) + p - n - 1$.

In fact, these are the only redundant conditions. Anderson [2] shows the following, in part using a geometric version of the above argument.

Theorem 4.2. *The set $E'(w)$ is the unique minimal set satisfying both*

- *We have $(p, q) \in E'(w)$ if and only if $(2n + 2 - p, 2n - q) \in E'(w)$.*
- *For any $v \in B_n$, $v \leq w$ if and only if $r_v(p, q) \leq r_w(p, q)$ for all $(p, q) \in E'(w)$.*

Furthermore, Anderson [2, p.13] also gives for each pair of triples $\{(p, q, r), (2n + 2 - p, 2n - q, p - q - 1 + r)\}$ the minimal element

$$v(p, q, r) = v(2n + 2 - p, 2n - q, p - q - 1 + r) \in B_n$$

such that $r_{v(p, q, r)}(p, q) > r$, thus explicitly giving a way of calculating $\mathcal{E}(w)$ as

$$\mathcal{E}(w) = \{v(p, q, r) \mid (p, q) \in E'(w), r = r_w(p, q)\}.$$

This correspondence is described in Figure 4 for the case where either $q < n$ or both $q = n$ and $p \geq n + 1$. Only the first half (meaning $w(1) \cdots w(n)$) of the elements $v(p, q, r)$ are listed, and $a \cdots b$ should be taken to be empty if $a > b$. The remaining cases can be inferred from the symmetry $v(p, q, r) = v(2n + 2 - p, 2n - q, p - q - 1 + r)$.

Consulting the table and calculating reduced expressions gives the following proposition.

Proposition 4.3. *Let $w \in B_n \subseteq S_{2n}$. Then w is defined by pseudo-inclusions if and only if, for all $(p, q) \in E(w)$, $v(p, q, r_w(p, q))$ has a unique reduced expression.*

Proof. By the symmetry of $E(w)$, we only need to consider the case where $q < n$ or both $q = n$ and $p > n$. If $q \geq p$, so $\max(0, q - p + 1) = q - p + 1$, then we have $q < n$ and hence $p + r \leq n$ by our assumptions, so

$$v(p, q, q - p + 1) = 1 \cdots (p - 2)p \cdots (q + 1)(p - 1)(q + 2) \cdots n = s_{n-p+1} \cdots s_{n-q},$$

where the indices in the last expression are decreasing.

Otherwise, we first treat the case $r = 0$. If $p \leq n$, then

$$v(p, q, 0) = 1 \cdots (q-1)pq \cdots (p-1)(p+1) \cdots n = s_{n-p+1} \cdots s_{n-q},$$

where the indices in the last expression are increasing. If $p > n$, then either $p+q < 2n+2$, in which case

$$v(p, q, 0) = 1 \cdots (q-1)pq \cdots (2n-p)(2n+2-p) \cdots n = s_{p-n-1} \cdots s_0 \cdots s_{n-q},$$

or $p+q \geq 2n+2$, in which case

$$v(p, q, 0) = 1 \cdots (2n-p)(2n+2-p) \cdots qp(q+1) \cdots n = s_{p-n-1} \cdots s_0 \cdots s_{n-q}.$$

In both cases, we take indices in the last expression to be first decreasing then increasing.

Finally, if $p = n+1$, $q = n$, and $r = 1$, then

$$v(n+1, n, 1) = 1 \cdots (n-2)(n+1)(n+2) = s_0 s_1 s_0.$$

These are the only elements of B_n with a unique reduced expression, so the proposition is proved. \square

Unfortunately, the proposition applies to $E(w)$ and not $E'(w)$, so it does *not* imply a statement about the Coxeter-theoretic coessential set $\mathcal{E}(w)$. The following example makes this difference clear.

Example 4.1. Let $n = 3$ and $w = 426153$. Then $E(w) = \{(3, 2), (5, 2), (5, 4), (3, 4)\}$. We see that w is not defined by pseudo-inclusions since $r_w(3, 2) = 1 \neq \max(0, 2 - 3 + 1)$. Moreover, $v(3, 2, 1) = v(5, 4, 1) = 351624 = s_1 s_0 s_2 s_1 = s_1 s_2 s_0 s_1$. However, $E'(w) = \{(5, 2), (3, 4)\}$, with $r_w(5, 2) = 0$. The permutation $v(5, 2, 0) = v(3, 4, 2) = 153426 = s_1 s_0 s_1$, which does have a unique reduced expression. Note $153426 < 351624$ and $\mathcal{E}(w) = \{153426\}$. The element w fails the Hultman condition since, if $u = 132546$, then $\ell_D(u, w) = 4$ but $\ell_T(u, w) = 2$. It may be significant that this element also fails to satisfy Proposition 2.6, as noted at the end of Section 2.

ACKNOWLEDGMENTS

I thank William Slofstra for helpful discussions. This paper was completed while I am on sabbatical at the Department of Mathematics at the University of Illinois at Urbana-Champaign, and I thank the department for its hospitality. I am supported by Simons Collaboration Grant 359792.

REFERENCES

- [1] Michael H. Albert and Robert Brignall, *Enumerating indices of Schubert varieties defined by inclusions*. J. Combin. Theory Ser. A **123** (2014), 154–168.
- [2] David Anderson, *Diagrams and essential sets for signed permutations*. Preprint available as arXiv:math/1612.08670.
- [3] Ron M. Adin, Christos A. Athanasiadis, Sergi Elizalde, and Yuval Roichman, *Character formulas and descents for the hyperoctahedral group*, Adv. Appl. Math. **87** (2017), 128–169.
- [4] Sara Billey and Tom Braden, *Lower bounds for Kazhdan–Lusztig polynomials from patterns*. Transform. Groups **8** (2003), 321–332.

- [5] Sara Billey and Alexander Postnikov, *Smoothness of Schubert varieties via patterns in root systems*. Adv. Appl. Math. **34** (2005), 447–466.
- [6] Anders Björner and Francesco Brenti, *Combinatorics of Coxeter groups*. Graduate Texts in Mathematics **231**, Springer, New York, 2005.
- [7] Anders Skovsted Buch, Andrew Kresch, and Harry Tamvakis, *A Giambelli formula for isotropic Grassmannians*. Selecta Math. (N.S.) **23** (2017), 869–914.
- [8] R. W. Carter, *Conjugacy classes in the Weyl group*. Compositio Math. **25** (1972), 1–59.
- [9] William Y. C. Chen, George Z. Gong, and Jeremy J. F. Guo, *The sorting index and permutation codes*. Adv. Appl. Math. **50** (2013), 367–389.
- [10] Matthew J. Dyer, *On minimal lengths of expressions of Coxeter group elements as products of reflections*. Proc. Amer. Math. Soc. **129** (2001), 2591–2595.
- [11] William Fulton, *Flags, Schubert polynomials, degeneracy loci, and determinantal formulas*. Duke Math. J. **65** (1992), 381–420.
- [12] V. Gasharov and V. Reiner, *Cohomology of smooth Schubert varieties in partial flag manifolds*. J. London Math. Soc. (2) **66** (2002), 550–562.
- [13] Meinolf Geck and Sungsoo Kim, *Bases for the Bruhat-Chevalley order on all finite Coxeter groups*. J. Algebra **197** (1997), 278–310.
- [14] Thomas Gobet, *On cycle decompositions in Coxeter groups*, Sémin Lothar. Combin. **78B** (2017), FPSAC 2017, Art. 45, 12 pp.
- [15] Axel Hultman, Svante Linusson, John Shareshian, and Jonas Sjöstrand, *From Bruhat intervals to intersection lattices and a conjecture of Postnikov*. J. Combin. Theory Ser. A **116** (2009), 564–580.
- [16] Axel Hultman, *Inversion arrangements and Bruhat intervals*. J. Combin. Theory Ser. A **118** (2011), 1897–1906.
- [17] Myrto Kallipoliti, *The absolute order on the hyperoctahedral group*. J. Algebraic Combin. **34** (2011), 183–211.
- [18] Richard Kane, *Reflection groups and invariant theory*. CMS Books in Mathematics/Ouvrages de Mathématiques de la SMC **5**, Springer–Verlag, New York, 2001.
- [19] Alain Lascoux and Marcel-Paul Schützenberger, *Treillis et bases des groupes de Coxeter*. Elect. J. Combin. **3(2)** (1996) Research Paper 27, approx. 35 pp.
- [20] Joel Brewster Lewis and Alejandro H. Morales, *Combinatorics of diagrams of permutations*, J. Combin. Theory Ser. A **137** (2016), 273–306.
- [21] Alexander Postnikov, *Total positivity, Grassmannians, and networks*. Preprint available as arXiv:math/0609764
- [22] Victor Reiner, *Signed permutation statistics and cycle type*. European J. Combin. **14** (1993), 569–579.
- [23] Victor Reiner, Alexander Woo, and Alexander Yong, *Presenting the cohomology of a Schubert variety*. Trans. Amer. Math. Soc. **363** (2011), 521–543.
- [24] Edward Richmond and William Slofstra, *Staircase diagrams and enumeration of smooth Schubert varieties*. J. Combin. Theory Ser. A **150** (2017), 328–376.
- [25] Jonas Sjöstrand, *Bruhat intervals as rooks on skew Ferrers boards*. J. Combin. Theory Ser. A **114** (2007), 1182–1198.
- [26] Henning Ulfarsson and Alexander Woo, *Which Schubert varieties are local complete intersections?* Proc. London Math. Soc. (3) **107** (2013), 1004–1052.

DEPT. OF MATHEMATICS, UNIVERSITY OF IDAHO, MOSCOW, ID 83844-1103

E-mail address: awoo@uidaho.edu